

Section 7 (Stat 171)

- TF: Jiaze Qiu
 - Email: jiazeqiug.harvard.edu
 - Section/OH: Wed 9-12 pm (ET)
- TF: Fernando Vicente
 - Email: fernando_vicente@fas.harvard.edu
 - Section/OH: Mon 7-9 pm and Thu 8-10 pm (ET)
- Acknowledgement: This handout is partially based on notes created by Lisa Ruan and Christy Huo.
- All the section materials (handouts & solutions) can be found either on Canvas or [here](#).

1 Poisson Processes Review

- A **counting process** $(N_t)_{t \geq 0}$ is a stochastic process such that:
 - $N_t \geq 0$ (non-negative)
 - N_t is an integer
 - $s \leq t \rightarrow N_s \leq N_t$ (non-decreasing)
- A **Poisson process** with parameter λ (not to be confused with poisson random variable) is a special type of counting process that further satisfies:
 - $N_0 = 0$
 - $N_t \sim \text{Pois}(\lambda t)$
 - For $s, t > 0, N_{t+s} - N_s \sim N_t \sim \text{Pois}(\lambda t)$
- A poisson process can also be characterized using exponential random variables:

$$N_t = \max \{n : X_1 + X_2 + \dots + X_n \leq t\}, \quad X_1, X_2, \dots \sim \text{Exp}(\lambda)$$

This implies:

$$X_k = S_k - S_{k-1}, \quad S_k = X_1 + X_2 + \dots$$

Where S_k also denotes the kth arrival time in a Poisson process.

- **Thinning/Superposition:** Both the thinned and superpositioned Poisson processes are also Poisson processes.
- **Conditional Poisson Process:** In addition to being related to exponential random variables, Poisson processes are also related to uniform random variables in the form of order statistics. Let S_1, S_2, \dots, S_n be the arrival times of a Poisson process:
 - $(S_1, S_2, \dots, S_n) \mid N_t = n \sim$ order statistics on $\text{Unif}[0, t]$
 - $f_{(S_1, S_2, \dots, S_n) \mid N_t = n}(S_1, S_2, \dots, S_n) = \frac{n!}{t^n}$ where $0 \leq S_1 \leq \dots \leq S_n$
- **Spatial Poisson Process:** A spatial Poisson process $(N_A)_{A \subseteq \mathbb{R}^d}$ with parameter λ satisfies the following:
 - For each bounded set $A \subseteq \mathbb{R}^d, N_A \sim \text{Pois}(\lambda|A|)$
 - If A and B are disjoint, N_A is independent from N_B

Note that a spatial Poisson process is essentially a Poisson process generalized to higher dimensions.

2 Practice problems

2.1 Midterm 1 Problem 4(b)

Suppose a Markov chain X_0, X_1, \dots has the transition matrix

$$P = \begin{pmatrix} 1/3 & 1/3 & 1/3 & 0 \\ 1/3 & 1/3 & 0 & 1/3 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad (1)$$

Given that the chain starts in state 1 and is eventually absorbed into state 3, what is the expected number of steps until the chain is absorbed.

2.2 Spatial Poisson

Consider a spatial Poisson process with parameter λ in \mathbb{R}^3 . Find the distribution of the distance from the origin to the nearest point/event. How is it related to the exponential distribution?

2.3 Conditional Poisson

Let Λ be a positive random variable with PDF p . Let $(N_t)_{t \geq 0}$ be a counting process, such that given $\Lambda = \lambda$, $(N_t)_{t \geq 0}$ is a Poisson process with rate λ .

1. Find the PMF for N_t .

2. If for all $s, t > 0$, $N_{t+s} - N_s \sim N_t$, our process is said to have stationary increments. If for any $0 \leq q < r \leq s < t$, $N_t - N_s$ is independent of $N_r - N_q$, it's said to have independent increments. Show that the our process has stationary increments, and briefly explain why it doesn't have independent increments in general.